

# Triple superphosphate, potassium sulfate and nitrogenous fertilizers effects on fitness and aggressiveness of *Fusarium culmorum* inducing wheat crown rot

Siham BAHA EDDINE

Laboratoire de biotechnologie botanique et de protection des végétaux, Département de biologie, Faculté des sciences, Université Ibn Tofail, Kenitra, Maroc

Allal DOUIRA

Laboratoire de biotechnologie botanique et de protection des végétaux, Département de biologie, Faculté des sciences, Université Ibn Tofail, Kenitra, Maroc

Brahim EL YOUSFI

Laboratoire de phytopathologie des céréales, Institut National de la Recherche Agronomique, Centre Régional de Settat, Maroc

This study examined the effects of N, P, and K fertilizers on *Fusarium culmorum* on in-vitro mycelial growth and biomass, and aggressiveness on wheat plants. Urea, phosphorus, potassium sulfate, and the mixture of similar amounts of these last two ingredients increased fungal biomass. Rearing mycelia on urea, the mixture, or phosphorus increased aggressiveness by 56%, 120%, and 130%, respectively. But this aggressiveness did not affect inoculated plants dry biomass. However, the inoculum that was reared on ammonium sulfate increased infected plant biomass. These findings infer that the management of this disease may rely on a proper dose of fertilizers application. First, farmers are advised to avoid urea and use instead ammonium nitrate; second, place phosphorus and potassium sulfate under the seedbed away from any inoculum in the soil. Therefore, surveying disease development, its past events, soil health, and soil fertility is a prerequisite for any successful control of this disease.

**Keywords:** *Fusarium culmorum*, Aggressiveness, Fitness, Nitrogen forms, Phosphorus, Potassium sulfate

## INTRODUCTION

Worldwide, *Fusarium culmorum* (*F. culmorum*) (W.G. Smith Sacc.) induces wheat crown rot (WCR) disease in arid and semi-arid regions (Burgess et al., 2001; Pettitt et al., 2003). This pathogen dominates the etiology of this disease, and especially within spring cereal crops (Wallace, 1978; Smiley et al., 2016; Voigt, 2002), while in other areas it is *F. Pseudograminearum* that is the causative agent of crown rot (Alahmad et al., 2018; Kazan and Gardiner, 2018). Gargouri et al. (2003) gave evidence that the genetic variability of this pathogen has no geographic origin. However, this variability only occurs within the pathogen's population and stems from large spore dispersal (Mishra et al., 2003).

Weather and crop management tools such as crop type, tillage, rotation, soil fertility, weather conditions, and their interactions highly affected pathogen infection and WCR disease development (Colhoun et al., 1968. Scherm et al., 2013; Smiley, 2016). Humidity and temperature affected mycelia biomass, and pathogenicity, and consequently contributed to the development of this disease (Cook and Christen, 1976; Brennan et al., 2003).

In water-stressed environments, infected cereal plants develop severe disease symptoms in comparison to those grown under irrigated conditions (Smiley et al., 1996; Chekali et al., 2011). Also, any cause that impels a decreasing state in soil water potential predisposes small grain cereals to this disease by increasing *F. culmorum* colonization of wheat seedlings (Papendick and Cook, 1974; Beddis and Burgess, 1992).

For disease control, scientists mostly look for a decrease of soil inoculum, through crop residue management and rational use of nitrogenous fertilizers (Cook, 1974; Chakraborty et al., 2006). Swan et al. (2000) promoted residue decomposition practices that mitigate disease incidence in the following crop through lowering *F. culmorum* survival (Warren and Kommedahl, 1973). In this later work, nitrogen application, in the form of ammonium nitrate, speeded wheat residues decomposition and thus reduced populations of *Fusarium* spp. (Warren and Kommedahl, 1973; Pereyra et al., 2004). Also, applying *Trichoderma harzianum* and nitrogen to infected plant residues displaced the pathogen from its niche, but this displacement depended on the form of nitrogen applied (Lakhesar et al., 2010). However, the interaction of the host and the pathogen in disease development is modulated by environmental factors and crop management tools (Voigt, 2002).

Most studies that dealt with nitrogen effect on disease development pointed out that overuse of nitrogenous fertilizers intensified the disease (Papendick and Cook, 1974; Rowaished, 1981; Kommedahl, 1984). Furthermore, the form of nitrogen has a greater effect on host resistance or disease severity than does its amount. But, with no specification of nitrogen form used, Hemissi et al. (2018) pointed out that nitrogen applied at 150 to 200 kg ha<sup>-1</sup> increased disease severity and whiteheads development. In contrast, urea and ammonium sulfate increased WCR severity, when they were applied as a baseline, or as top-dressing. But, ammonium nitrate application, at pre-sowing or at tillering, and/or at stem elongation, significantly reduced the disease (Baha Eddine et al., 2019).

To our knowledge, limited studies have dealt with pathogen fitness and aggressiveness under nitrogen forms and/or other macronutrients action (Garrett, 1976; Huber, 1990). In any pathosystem (Huber and Haneklaus, 2007), nutrients play a greater role in altering the pathogen's virulence (Thomas and Elkinton, 2004). Therefore, the nutrients' effects on the pathogen biology and the host, through temporal interaction among disease components are a prerequisite to the control of this disease (Agrios, 2005; Chakraborty et al., 2006). The general use of fertilizers (N, P, and K) in cereal production (FAO, 2006) gave a reason for this study to test them as treatment factors. Therefore, we examined the effect of nitrogenous, phosphorus, and potassium fertilizers on *F. culmorum* fitness and aggressiveness.

## MATERIALS AND METHODS

### Fertilizers' effect on in-vitro mycelium growth and biomass

This experiment used phosphorus as triple superphosphate (45% P), potassium as potassium sulfate (48% K), phosphorus plus potassium sulfate (1/2-1/2 for each), and nitrogen as ammonium sulfate (21% N), ammonium nitrate (33.5% N) and urea (46% N) as fertilizers. From each fertilizer, six quantities were independently added to flasks that contained one liter of distilled water and 10 g of glucose. These quantities were rated 1 as control (unfertilized treatment), and 2: 0.2 g, 3: 0.4 g, 4: 0.8 g, 5: 1.2 g, and 6: 1.6 g as the remaining levels. All the flasks were sterilized for 20 minutes at 120 °C and a pressure of 3 bars. Once cooled, each flask was inoculated with a PDA fragment (approximately 1 cm<sup>2</sup>) of a fresh local virulent *F. culmorum* isolate. Note that *fusarium* isolation and identification followed the procedure described in our previous article (Baha Eddine et al., 2019). The isolate used herein was obtained from infected crown wheat. Inoculated flasks were incubated for one month under 12h of photoperiod. Afterward, the content of each flask was separately filtered through sterilized cheesecloth, and the collected mycelia biomass was independently dried out in a laminar flow cabinet, and then weighed.

This experiment was repeated once. Note that the flasks of the first experiment were incubated at 20–25 °C, whereas those of the second one were incubated at 15–20 °C. Additionally, for the first trial, a mycelium sample from each treatment (fertilizer by dose interaction) was transferred to two Petri dishes containing PDA medium. This fungal transfer is meant to assess the carryover effect of fertilizers and their rates on in- vitro mycelia growth.

Colony growth measurements in each Petri dish were initiated on the 3rd day and resumed on the sixth day of post-inoculation. The evolution of the growth colony (cm), in four perpendicular directions of a plate, was monitored. The colony size was estimated by the following formula:

Where  $R_i$  is the radial length in direction  $i$ , and the mean of colony size per treatment by dose and by experiment was plotted against post-inoculation dates (Baha Eddine et al., 2020). This monitoring of the growth is intended to assess the fungal fitness (Pringle and Taylor, 2002).

### **Fertilizers' effect on Aggressiveness**

The experiment was carried out in a greenhouse in the 2017–2018 growing season, and was laid out in the first week of November and ended in late April. Six plastic trays of 77 cavities (7 × 11) were used and each cavity had a dimension of 64 cm<sup>3</sup> (4 × 4 × 4 cm). Each tray contained one fertilizers' treatment level: phosphorus, potassium sulfate, phosphorus plus potassium sulfate, ammonium sulfate, ammonium nitrate, or urea. Within each plastic tray, six columns that were vertically arranged represented fertilizers' rates, each of which was separated by a column of 7 voids. And within each fertilizer's column, the seven cavities represented repetitions (mycelium samples as batches). All the cavities, except those separating the rate's factor, were filled to half of their capacity with a natural soil (Tirs) brought from the experimental station of Sidi El Aïdi (INRA, Settat, Morocco). This soil was previously sieved and sterilized, for one hour, three times in a row at 120 °C and a pressure of 3 bars. Five seeds of a sensitive durum wheat cultivar Ourgh were sown in each cavity, and then a thin layer of the same soil was broadcast to cover the seeds. Thereafter, 12 ml of water was poured onto each seeded cavity to initiate germination.

Two days later, the dried mycelia biomass, collected from all liquid media, were separately grounded and homogeneously distributed over the 7 vertical cavities which represented batches. The inoculum was then covered with a thin layer of sterile soil. Irrigation with distilled water was provided according to the plant's need.

Pathogen aggressiveness defined as the pathogen's ability to invade and establish in the host (Thomas and Elkinton, 2004) was quantified herein by the severity of symptoms induced by the inoculum reared on different growth media. Disease evaluation started at the heading growth stage in line with our previous work (Baha Eddine et al., 2019). Any plant with a disease progression that exceeded the third node had been given a score of 3 as the highest severity score. Subsequently, evaluated plants per treatment were separately left to dry in the open air of a shutdown greenhouse. Then, the dry weight of plant biomass affected by each treatment was weighted. Note that this experiment was immediately carried out after each trial that dealt with mycelia biomass production. This experiment was repeated once.

### **Statistical Analysis**

All statistical computations were carried out using the SPSS statistical package (version 20) for Windows. A combined analysis, of the two experiments, was used to assess treatment effect (fertilizers and rates) on fungal biomass. The Leven test for homogeneity of variance between the two experiments gave a basis to separate analysis for each experiment. Therefore, fertilizers' effect and their doses, within each experiment, were evaluated by an analysis of variance (ANOVA) using the GLM procedure. Significant differences among fertilizers were tested at 5% probability with the Waller-Duncan test statement, and doses with a polynomial contrast statement (West et al., 2015). And to analyze the strength of this relationship either linear or logarithmic, a regression analysis

was performed. On the other hand, to evaluate the carryover effect, a mixed model with colony size as a dependent variable was adopted with fertilizers and rates as fixed factors and plates as random. The experimental design was a longitudinal model with a repeated statement as post-inoculation days and subjects as plates. We selected a covariance matrix as arima 1 with the covtype (ar1) statement and everything else is kept as default. The experiment about disease severity was analyzed as a nested design. Fertilizers were taken as a fixed factor while doses were nested within fertilizers because they were assumed to have different nitrogen concentrations and then were taken as experimental errors for fertilizers. The batches (samples of inoculum) within doses by fertilizers were experimental error for doses nested within fertilizers. Significant differences between the control and fertilizers' effect were based on Bonferroni test at 5% probability. Finally, the two levels of this later experiment were taken as random (Berger et al., 2018).

## RESULTS

The two trials about fungal biomass production had a highly significant different variance (Levene statistic = 7.09;  $df_1 = 1$ ;  $df_2 = 70$ ;  $P = 0.010$ ). This significant test was the basis for separately analyzing them. Therefore and within both experiments, fertilizers and their rate were found to significantly affect *F. culmorum* biomass ( $P \leq 0.001$ ). Also, the polynomial contrast test gave evidence that the increase of biomass production with an increase in fertilizers' rates had a significant cubic profile ( $P \leq 0.001$ ) in both trials.

The use of ammonium sulfate, ammonium nitrate, urea, or phosphorus as a substrate in growth media, induced a similar increase of fungal biomass, but this increase was different within the two temperature essays (Figures 1a, 1b, 1c, and 1d). In contrast, potassium sulfate, or the mixture of phosphorus, and potassium sulfate mostly produced similar biomass production profiles nearly smaller between the two experiments (Figures 1e and 1f). The significant effect of the growth media on fungal biomass showed that ammonium sulfate, ammonium nitrate, or potassium sulfate gave the lowest production in both experiments (Table 1). Whereas, those containing urea, phosphorus, or the phosphorus mixture and potassium sulfate significantly stimulated *F. culmorum* in-vitro growth, and therefore the fungal biomass (Table 1).

A significant relationship between fertilizers' doses and *F. culmorum* fungal biomass production is evident in Table 2, except for ammonium sulfate and ammonium nitrate at a temperature of 15 to 20 °C (Table 2). Most of the other doses had a logarithmic relation between doses and biomass production. But potassium doses showed a significant linear effect even stronger at 20 to 25 °C when comparing its standardized coefficient beta of 0.95 to 0.76 at 15-20 °C. Also, the use of urea, phosphorus, or the mixture of phosphorus and potassium induced similar logarithmic strength on the production of fungal biomass under both temperature conditions (Table 2).

Overall, the two experimental conditions had a different effect on fungal biomass (Figure 1), and within each trial, the main effect of fertilizers' rate level off at 0.2 g L<sup>-1</sup>.

The inoculum reared on different sources of fertilizers and rates presented similar growth patterns on PDA medium (Table 3), and the only significant effect ( $P = 0.001$ ) on in-vitro mycelial growth was relative to days of post-inoculation (Table 3). Therefore, the growth media continent did not have any carryover effect on the fitness cost of this pathogen.

Fertilizers significantly affected disease severity ( $P \leq 0.001$ ) and therefore the aggressiveness of the pathogen (Table 4). Compared with the unfertilized check, the overall use of the mycelia reared either on urea, the mixture of phosphorus and potassium sulfate, or phosphorus stimulated disease severity by 56%, 120%, and 130%, respectively (Figure 2). In contrast, amended media with ammonium sulfate and ammonium nitrate did not change this characteristic (Figure 2).

All fertilizers, rates within fertilizers, and batches significantly affected the dry weight of inoculated plants at  $P \leq 0.001$ ,  $P \leq 0.001$ , and  $P \leq 0.048$ , respectively (Table 5). The greatest inoculated plant dry weight was induced by the inoculum cultured on ammonium sulfate media (Figure 3). The significant effect of rates within fertilizers on infected plant biomass, depicted in Figure 4, revealed an interaction between fertilizers and their rates. This interaction was mostly fueled by differences between the effects of ammonium nitrate and ammonium sulfate (Figure 4).

## DISCUSSION

Levels of yield losses caused by WCR disease urge scientists and farmers to seek sustainable control measures based on the knowledge of disease biology and epidemiology (Duveiller et al., 2007; Paulitz et al., 2010; Chekali et al., 2013). Thus, we wanted to manage this disease with major mineral nutrients (nitrogen, phosphorus, and potassium), applied at different rates, on the fitness features (growth, biomass production, and aggressiveness) of *F. culmorum*.

This work was set up as a continuation of a preliminary study that dealt with *F. culmorum* growth on solid PDA (Damir, 2014). The solid media were independently amended with 0.1, 0.2, 0.4, and 0.8 g L<sup>-1</sup> of phosphoric, potash, and nitrogenous fertilizers. Though, media containing phosphorus at 0.4 g L<sup>-1</sup>, 0.6 g L<sup>-1</sup>, and 0.8 g L<sup>-1</sup> remained liquid, although we increased the amount of agar. And these findings justified our use of liquid media instead.

Monitoring *F. culmorum* biomass production, within the two-temperature conditions, showed that the mycelia production was dependent on the setup of the two experiments. The fungal biomass obtained at 20-25 °C was greater than the one obtained at 15-20 °C by 104% (Table 1). This finding is supported by the optimum growth at 25 °C of the European *Fusarium culmorum* isolates on PDA, irrespective of their geographic origin, and these later isolates were the fastest-growing of the fusaria tested (Brennan et al., 2003). Under these two temperature conditions, fertilizers affected the fitness and aggressiveness of *F. culmorum*. Phosphorus and the mixture (phosphorus and potassium sulfate) improved these two features, and explain in part Greaney's (1938) observation at Winnipeg who noticed an increase in disease severity when soil phosphorus was in excess or a deficient state. Damir (2014) also pointed out that phosphorus added at post-emergence to inoculated plants, grown in pots filled with natural soil, enhanced disease development. Further, she found that the application of phosphorus and potassium sulfate as a solution at post-emergence increased disease severity on the same variety herein used. These reported studies infer that close contact of this pathogen with phosphorus acts directly upon the fitness components, and therefore impacted disease development.

Even though the application of phosphorus and its addition to potassium sulfate increased the fitness, aggressiveness, and disease development, they did not affect the infected plant biomass (Table 1, 4, and 5, Figures 2 and 3). Baha Eddine et al. (2019) clearly showed that WCR disease development has a close association with nitrogen fertilizers. Thus, whenever infected plants were fertilizers with either form of ammonium sulfate or urea, the disease intensity as the incidence and severity took severe proportions. In this study, we did not add any fertilizer to the natural soil, and levels of disease severity recorded may infer that the disease stems from a nitrogen pool (mycotoxins promoters) of the pathogen. Therefore, adding nitrogen shifted this pathogen from a non-dominant to a host-dominant disease in this pathosystem (Kommedahl and Windels, 1979).

Although potassium sulfate had shown some effect on fitness and aggressiveness it did not induce any significant disease severity (Table 1 and Figure 2). Hofgaard et al. (2010) also noticed a reduction in-vitro growth of *F. culmorum* and *F. graminearum* when the fertilizer containing potassium phosphate was added into the growth medium. Even a decrease in head blight was observed when this fertilizer was applied as a foliar application on wheat plants. Also, Onuorah (1969) gave evidence that an application of potassium sulfate decreased WCR severity on adult plants. Moreover, high potassium application, especially if associated with high nitrogen, reduced



the disease even further (Dixon et al., 2010).

Application of a  $\text{CaNO}_3$  ( $1.21 \text{ g L}^{-1}$ ) solution on infected plants increased WCR disease development (Onuorah, 1969). And Papendick and Cook (1974) further demonstrated that application of nitrogen when coupled to water stress enhanced disease incidence and severity. But Rowaished (1981) used a series of Hoagland solutions containing concentrations and forms ( $\text{NO}_3$  and  $\text{NH}_4$ ), ranging from 21 ppm up to 1050 ppm, and found a slight increase in disease severity with high concentration levels of nitrogen.

At the two temperatures, nitrogen, like urea, affected fungal biomass, aggressiveness, and overstimulated this biomass by one to two folds (Table 1). This significant increase induced levels of aggressiveness that were less than those of phosphorus and the mixture but did not affect plant dry biomass. In recent work, urea was also found to optimally increase pathogen fitness when it was added to solid media at a rate of  $1.5 \text{ g L}^{-1}$  (Baha Eddine et al., 2020). But concentrations over  $12 \text{ g L}^{-1}$  of urea were harmful to the pathogen. However, at a fixed nitrogen concentration of 1.57 % N among nitrogenous fertilizers, this nutrient favored mycelium growth and aggressiveness by 50%. Additionally, Damir (2014) also showed that an application of urea either at  $0.1$  or at  $0.2 \text{ g L}^{-1}$  favored fungal growth. This nutrient was also found to increase disease development when placed either under seedbed or as a foliar application (Baha Eddine et al., 2019), or as a top-dressing fertilizer (Akgül and Erkilic, 2016; Baha Eddine et al., 2019).

Sulfur is known to be a good fungicide against various plant diseases (Haneklaus, 2007). But surprisingly in our study, ammonium sulfate was found to only affect growth fitness at  $25^\circ\text{C}$ , and on the contrary, increased dry weight of infected plants through a non-significant effect on disease severity. In another work, fertilizing infected plants, independently as top-dressing at tillering or at stem elongation, induced disease severity by 33% and 30%, respectively (Baha Eddine et al., 2019), but nothing was said on yield. Yet, a simultaneous application of this fertilizer at these two growth stages, increased disease severity by 53% (Baha Eddine et al., 2019). Ammonium sulfate with 23 to 24% of sulfate also has a fungistatic action on wheat powdery mildew (Cook, 1987). Also, Williams and Cooper (2004) gave evidence that some fusaria species were sensitive to this nutrient. This may explain in our work the noted weak fitness and aggressiveness of this pathogen, but its stimulation of the infected plant biomass needs more research.

The mitigated ammonium nitrate's effect on fitness (Table 1) and aggressiveness (Figure 2) agreed with the findings of Baha Eddine et al. (2020) which stated that the use of this fertilizer at  $24 \text{ g L}^{-1}$  lessened these two characteristics, and resulted in a decrease of disease severity with a reliable expression of disease resistance. Thus, these results explain why levels of this disease were low when this fertilizer was applied at pre-sowing or as a top dressing at tillering and/or at stem elongation (Baha Eddine et al., 2019).

The macronutrient impact on plant diseases depends on host response, previous crop sequence, rate of nitrogen applied, residual N, the period of application, soil microorganisms, the ratio of  $\text{NH}_4^+$  to  $\text{NO}_3^-$  and disease etiology (Kommedahl, 1984). However, depending on the pathosystem, other companion ions such as Ca, K, Cl, and  $\text{SO}_4$  may also be involved in disease development (Dordas, 2008; Lawrence and Wade, 2018, Yadav et al., 2020). Thus, every pathosystem must be investigated for the right nutrient forms to use for any disease management (Fageria et al., 2010). Evaluating the effect of N, P, K nutrients on this fungus may shed light on its genetic diversity characterized by phenotypic traits such as pathogenicity, aggressiveness, and mycotoxin production (Korn et al., 2011).

## CONCLUSION

This study added new evidence to the causal relation between nitrogen form and disease development. Therefore, it provided insight into WCR disease by assessing the interaction between

the causal agent and N, P, and K macronutrients. The use of urea and phosphorus alone or with potassium sulfate increased *F. culmorum* fitness and aggressiveness. In contrast, ammonium nitrate, as a nutrient, significantly lessened these two characteristics. The study also allowed us to assume that these findings may pinpoint some disease management, and therefore, fulfill the control of this disease throughout soil health quality. For example, when *F. culmorum* propagules occur in soil profiles, we can speculate that the application of urea for top-dressing cereal in arid and semi-arid regions may not be recommended, whereas the use of ammonium nitrate can be a good alternative. Also, we advise cereal farmers to incorporate, into their field, phosphorus, and potassium under seedbed or drilled rowed to avoid any contact of these fertilizers with *F. culmorum*, and decline any furrow application. However, the use of ammonium sulfate as a baseline fertilizer is recommended to take advantage of this fertilizer's fungistatic action. But preliminary tests for soil fertility and quantification of the inoculum in cereal fields at pre-sowing frame valuable information for crop management in regions prone to WCR disease. Finally, scientists involved in screening and quantifying disease resistance may consider these findings when carrying out their selection methods. Other scientists may focus on the nutrition effect on the molecular and epigenetic status of *F. culmorum*. For functional annotation and confidence in our finding, confirmation of our results under field conditions is deemed necessary.

## REFERENCES

- Agrios G.N. (1997). *Plant Pathology*, 4th eds. Academic Press, New York (US).
- Alahmad S., Simpfendorfer S., Bentley A.R., Hickey L.T. (2018). Crown rot of wheat in Australia: *Fusarium pseudograminearum* taxonomy, population biology, and disease management. *Australasian Plant Pathology*, 47: 285-99.
- Akgül D.S., Erkilic A. (2016). Effect of wheat cultivars, fertilizers, and fungicides on *Fusarium* foot rot disease of wheat. *Turkish Journal of Agriculture and Forestry*, 40: 101-108.
- Baha Eddine S., El Yousfi B., Douira A. (2019). Interaction of nitrogen fertilizers with wheat growth stage and foliar treatment with urea effects on WCR induced by *Fusarium culmorum*. *Plant Archives*, 19: 2829-2835.
- Baha Eddine S., El Yousfi B., Douira A. (2020). Effects of nitrogen forms and rates on *Fusarium culmorum* growth, fitness, aggressiveness, and wheat, barley, and triticale resistance to crown rot disease. *Plant Cell Biotechnology and molecular biology*, 21: 107-129.
- Beddis A., Burgess W. (1992). The influence of plant water stress on infection and colonization of wheat seedlings by *Fusarium graminearum* group 1. *Phytopathology*, 82: 78-82.
- Berger P.D., Maurer R.E., Celli G.B. (2018). *Experimental design with applications in Management, Engineering, and the Sciences*. Second Edition. Springer International Publishing Company.
- Brennan J.M., Fagan B., Van Maanen A., Cooke B.M., Doohan F.M. (2003). Studies on in vitro growth and pathogenicity of European *Fusarium* fungi. *European Journal of Plant Pathology*, 109: 577-587.
- Burgess L.W., Backhouse D., Summerell B. (2001). Crown rot of wheat. In *Fusarium: Paul E. Nelson Memorial Symposium* (Eds. Summerell BA, Leslie IF, Backhouse D, Bryden WL & Burgess LW), pp. 271-294. The APS Press, Minnesota (US).
- Chakraborty S., Liu C.J., Mitter V., Scott J.B., Akinsanmi O.A., Ali S., Dill-Macky R., Nicol J., Backhouse D., Simpfendorfer S. (2006). Pathogen population structure and epidemiology are keys to WCR and *Fusarium* head blight management. *Australasian Plant Pathology*, 35: 643-655.

- Chekali S., Gargouri S., Berraies S., Gharbi M.S., Nicol M.J., Nasraoui B. (2013). Impact of Fusarium foot and root rot on yield of cereals in Tunisia. *Tunisian Journal of Plant Protection*, 8: 75-86.
- Chekali S., Gargouri S., Paulitz T., Nicol J.M., Rezgui M., Nasraoui B. (2011). Effects of Fusarium culmorum and water stress on durum wheat in Tunisia. *Crop Protection*, 30: 718-725.
- Colhoun J., Taylor G.S., Tomlinson R. (1968). Fusarium diseases of cereals: II. Infection of seedlings by *F. culmorum* and *F. avenaceum* in relation to environmental factors. *Transactions of the British Mycological Society*, 51: 397-404.
- Cook R.J., Christen A.A. (1976). Growth of cereal root-rot fungi as affected by temperature-water potential interactions. *Phytopathology*, 66: 193-197.
- Cook R.J. (1974). Control of Fusarium foot rot of wheat in the Northwest USA. *Australasian Plant Pathology*, 3: 50.
- Cook R.J. (1987). Sulphur as a cereal fungicide. In *Proceedings of the International Symposium Elemental Sulphur in Agriculture: Nice, France, 25-27 March 1987*, 2: 417-424.
- Damir A. (2014). Effet de la fertilisation NPK sur la sévérité des pourritures racinaires, causées par *Fusarium culmorum* et *Bipolaris sorokiniana*, ainsi que son effet sur le rendement. *Projet de fin d'étude*. Université Hassan premier, Settat, Maroc.
- Dixon G.R., Tilston E.L. (2010). Soil-borne pathogens and their interactions with the soil environment. In *Soil microbiology and sustainable crop production*, pp. 197-271. Springer, Dordrecht.
- Dordas C. (2008). Role of nutrients in controlling plant diseases in sustainable agriculture. A review. *Agronomy for sustainable development*, 28: 33-46.
- Duveiller E., Singh R.P., Nicol J.M. (2007). The challenges of maintaining wheat productivity: pests, diseases, and potential epidemics. *Euphytica*, 157: 417-430.
- Fageria N.K., Baligar V.C., Jones C.A. (2010). *Growth and mineral nutrition of field crops*, 3rd ed. CRC Press, Taylor and Francis Group (US).
- FAO (2006). *Utilisation des engrais par culture au Maroc*. Première Ed. FAO, Rome.
- Gargouri S., Bernier L., Hajlaoui M.R., Marrakchi M. (2003). Genetic variability and population structure of the wheat foot rot fungus, *Fusarium culmorum*, in Tunisia. *European journal of plant pathology*, 109: 807-815.
- Garrett S.D. (1976). Influence of nitrogen on cellulolysis rate and saprophytic survival in soil of some cereal foot-rot fungi. *Soil Biology and Biochemistry*, 8: 229-234.
- Greaney F.J. (1938). The effect of phosphate deficiencies on infection of wheat by *Fusarium culmorum*. *Canadian Journal of Research*, 16: 27-37.
- Haneklaus S., Bloem E., Funder U., Schnug E. (2007). Effect of foliar-applied elemental sulphur on Fusarium infections in barley. *Landbauforschung Volkenrode*, 57: 213.
- Hemissi I., Gargouri S., Hlel D., Hachana A., Abdi N., Sifi B. (2018). Impact of nitrogen fertilization on Fusarium foot and root rot and yield of durum wheat. *Tunisian Journal of Plant Protection*, 13: 31-38.





- Hofgaard I.S., Ergon Å., Henriksen B., Tronsmo M. (2010). The effect of potential resistance inducers on development of *Microdochium majus* and *Fusarium culmorum* in winter wheat. *European journal of plant pathology*, 128: 269-281.
- Huber D.M., Haneklaus S. (2007). Managing nutrition to control plant disease. *Landbauforschung Volkenrode*, 57: 313.
- Huber D.M. (1990). Fertilizers and soilborne diseases. *Soil Use and Management*, 6: 168-172.
- Kazan K., Gardiner D.M. (2018). *Fusarium* crown rot caused by *Fusarium pseudograminearum* in cereal crops: recent progress and future prospects. *Molecular Plant Pathology*, 19: 1547-62.
- Kommedahl T. (1984). Interaction of nitrogen use and plant disease control. *Nitrogen in Crop Production* : 461-472.
- Kommedahl T., Windels C.E. (1979). Fungi: pathogen or host dominance in disease. *Developments in Agricultural and Managed Forest Ecology*, 5: 1-103.
- Korn U., Müller T., Ulrich A., Müller M.E.H. (2011). Impact of aggressiveness of *Fusarium graminearum* and *F. culmorum* isolates on yield parameters and mycotoxin production in wheat. *Mycotoxin research*, 27: 195-206.
- Lakhesar D.P.S., Backhouse D., Kristiansen P. (2010). Nutritional constraints on displacement of *Fusarium pseudograminearum* from cereal straw by antagonists. *Biological Control*, 55: 241-247.
- Lawrence E.D., Wade H.E. (2018). Mineral nutrition and florists' crops Diseases. In *Handbook of Florists' Crops Diseases* (Eds. McGovern R.J., Elmer W.H.), pp. 237-251. Springer.
- Mishra P.K., Fox R.T., Culham A. (2003). Inter-simple sequence repeats and aggressiveness analyses revealed high genetic diversity, recombination and long range dispersal in *Fusarium culmorum*. *Annals of Applied Biology*, 143: 291-301.
- Onuorah P.E. (1969). Effect of mineral nutrition on the *Fusarium* brown foot-rot of wheat. *Plant and soil*, 30: 99-104.
- Papendick R.I., Cook R.J. (1974). Plant water stress and development of *Fusarium* foot rot in wheat subjected to different cultural practices. *Phytopathology*, 64: 358-363.
- Paulitz T.C., Okubara P.A., Schroeder K.L. (2010). Integrated control of soilborne pathogens of wheat. In *Recent Developments in Management of Plant Diseases* (Eds. Gisi U, Chet I & Gullino ML), pp. 229-245. Springer, Dordrecht.
- Pereyra S.A., Dill-Macky R., Sims A.L. (2004). Survival and inoculum production of *Gibberella zeae* in wheat residue. *Plant Disease*, 88: 724-730.
- Pettitt T., Xiangming X., Parry D. (2003). Association of *Fusarium* species in the wheat stem rot complex. *European Journal of Plant Pathology*, 109: 769-774.
- Pringle A., Taylor J.W. (2002). The fitness of filamentous fungi. *Trends in microbiology*, 10: 474-481.
- Rowaished A.K. (1981). The influence of different forms of nitrogen on *Fusarium* root rot disease of winter wheat seedlings. *Journal of Phytopathology*, 100: 331-339.
- Scherm B., Balmas V., Spanu F., Pani G., Delogu G., Pasquali M., Migheli Q. (2013). *Fusarium culmorum*: causal agent of foot and root rot and head blight on wheat. *Molecular plant pathology*,



14: 323-341.

Smiley R.W., Collins H.P., Rasmussen P.E. (1996). Diseases of wheat in long-term agronomic experiments at Pendleton, Oregon. *Plant Disease*, 80: 813-820.

Smiley R.W., Machado S., Rhinhart K.E.L., Reardon C.L., Wuest S.B. (2016). Rapid quantification of soilborne pathogen communities in wheat-based long-term field experiments. *Plant Disease*, 100: 1692-1708.

Swan L.J., Backhouse D., Burgess L.W. (2000). Surface soil moisture and stubble management practice effects on the progress of infection of wheat by *Fusarium pseudograminearum*. *Australian Journal of Experimental Agriculture*, 40: 693-698.

Thomas S.R., Elkinton J.S. (2004). Pathogenicity and virulence. *Journal of invertebrate pathology*, 85: 146-151.

Voigt K. (2002). Management of *Fusarium* Diseases. In *Agricultural Applications*, pp. 217-242. Springer, Berlin, Heidelberg.

Wallace H.R. (1978). The diagnosis of plant diseases of complex etiology. *Annual Review of Phytopathology*, 16: 379-402.

Warren H.L., Kommedahl T. (1973). Fertilization and wheat refuse effects on *Fusarium* species associated with wheat roots in Minnesota. *Phytopathology*, 63: 103-108.

West B.T., Welch K.B., Gałecki A.T. (2015). *Linear mixed models, a practical guide using statistical software*, Second eds. CRC Press Taylor & Francis Group (US).

Williams J.S., Cooper R.M. (2004). The oldest fungicide and newest phytoalexin-a reappraisal of the fungitoxicity of elemental sulphur. *Plant Pathology*, 53: 263-279.

Yadav A.N., Mishra S., Kour D., Yadav N., Kumar A. (2020). *Agriculturally important fungi for sustainable agriculture. volume 1: perspective for diversity and crop productivity*. Springer, Cham.

## References